

Genetic Diversity in *Prunus spinosa* L. and Challenges in the Use of Autochthonous Sources

Klaus EIMERT

Geisenheim University, Center of Applied Biology, Institute of Botany, Germany

Article Info:

Research article

Corresponding author: Klaus EIMERT, e-mail: klaus.eimert@hs-gm.de

doi: <http://dx.doi.org/10.17474/acuofd.05189>

ABSTRACT

Genetic diversity in natural stands of autochthonous blackthorn (*Prunus spinosa* L.) of different German provenances has been analyzed using a highly reproducible high-annealing-temperature random amplified polymorphic DNA (HAT-RAPD) protocol. The findings were compared to those from seedstocks of the same provenances, reported earlier. Generally, genetic diversity in the natural stands was even lower (H_o 0.099–0.116) compared to the corresponding seedstocks (H_o 0.118–0.133). Furthermore, genetic differentiation was found to be moderate between natural residential sources (pairwise F_{st} 0.138–0.184, 22.527% variation among populations), but higher than between the seedstocks (pairwise F_{st} 0.086–0.104, 7.782% variation among populations). The findings are discussed in respect to German conservation law and its practical implementation.

Key words: *Prunus spinosa*; autochthonous; RAPD; genetic diversity; differentiation

Prunus Spinosa L.'nin Genetik Çeşitliliği ve Yerli Popülasyon Kaynaklarının Kullanımındaki Zorluklar

ÖZET

Almanya'nın farklı bölgelerindeki çakal eriği (*Prunus spinosa* L.) doğal türünün genetik çeşitliliği, yüksek yapışma sıcaklığında rastgele çoğaltılmış polimorfik DNA (HAT-RAPD) protokolü kullanılarak analiz edilmiştir. Bulgular daha önceki çalışmalarda kaydedilen benzer bölgelerden toplanmış tohum stoklarından elde edilen örneklerin bulgularıyla karşılaştırılmıştır. Genellikle doğal türlerdeki (suş) genetik çeşitlilik önceki çalışmalarda ölçülen değerden (H_o : 0,118-0,113) düşük olduğu (H_o : 0,099-0,116) belirlenmiştir. Dahası genetik farklılaşma doğal türlerde orta düzeyde (F_{st} ; 0,138- 0,184; popülasyonlar arası varyasyon oranı % 22,527) ama tohum stoklarından elde edilen örneklerin değerlerinden (F_{st} ; 0,086- 0,104; popülasyonlar arası varyasyon oranı % 7,782) daha yüksek olduğu belirlenmiştir. Bulgular Almanya koruma kanunları ve uygulamaları bakımından tartışılmıştır.

INTRODUCTION

Genetic diversity is generally agreed to be a main prerequisite for evolution providing raw material for selection (Rees et al. 2001, Crawford and Whitney 2010). Also natural genetic diversity provides raw material for breeding food and feed varieties (Hoisington et al. 1999, Esquinas-Alcázar

2005). While artificially increasing genetic diversity by mutation breeding or artificial crosses across breeding borders can increase genetic diversity, naturally occurring genetic diversity remains the largest pool of genes/allels to ensure healthy environments and secured food situation even during challenges like socio-ecological development and global climate

change. Thus, conservation of genetic diversity (by protection of environment) was agreed upon as a common goal for humanity (Convention on biological diversity 1992). Those agreements have to be implemented into national law by the subscribing countries. In Germany, the Federal Conservation Act was accordingly amended in 2010 (BnatSchG) to (among other points) include conservation of genetic diversity. Thus, it also regulates the introduction of non-resident species into the open landscape (outside settled areas and not used in agri-, horti-, or silviculture) - which will be prohibited from 2020 on. Until then there is a period of transition during which use of residential sources for any species is required whenever possible. In the BnatSchG, the term species is defined including sub-species levels like subspecies or even populations (BnatSchG 2010, §7(2)3). This regulation presents a practical problem for the use of plants in open landscape plantation. While an increasing number of publications are available for forest trees (Kremer et al. 2002, Petit et al. 2003, Magri et al. 2006) not much is known about genetic constitution of many other plants including shrubs, widely used in open landscape plantings. Despite this lack of knowledge, 9 regions of origin were introduced for woody plants (shrubs), not regulated by forestry or agricultural laws in 2003 (BMVEL 2003). Base for those regions were ecological basic units based on geographical classification of natural landscapes in Germany according to Schmidt and Krause (1997). This regulation does not take into account finer ecological structuring nor does it consider the different biologies of plants species. In 2012 the number of regions of origin was further reduced to 6 (BMU 2012).

Faced with this situation we aimed to broaden our knowledge on one of the species most commonly used for plantings

in the open landscape – blackthorn (*Prunus spinosa* L.).

Prunus spinosa L. (common blackthorn, sloe) is an insect-pollinated, animal-dispersed shrub native to Europe, North Africa, and West Asia (Schütt et al. 1992). It is very wide-spread over Germany and most of Europe, and therefore it is often used in open landscape plantation and renaturation measures in Germany. Blackthorn is assumed to be allo-tetraploid ($2n=4x=32$, Reynders-Aloisi and Grellet 1994) and it also propagates strongly by root suckers (Guitian et al. 1993).

In the recent years, several publications emerged dealing with the genetic structure of blackthorn. Mohanty et al. (2000, 2002) analyzed cp-DNA to observe large scale genetic structure in Europe, including a few scattered samples from Germany. Isozyme analyses for several areas in Germany showed moderate to low nuclear genetic diversity within and among blackthorn populations (Leinemann et al. 2002, Fronia 2009). Eimert et al. (2012) also reported relatively low genetic diversity in and no to weak differentiation among residential seedstock sources of blackthorn. Similar levels of genetic diversity were detected in eight autochthonous blackthorn populations from Flanders using AFLP and morphometry (Mijnsbrugge et al. 2013). Interestingly, in those populations, higher differentiation of nuclear DNA was reported, although no obvious morphometric differentiation was observed. Very recently, Leinemann et al. (2014) analyzed nuclear and chloroplast DNA of blackthorn from 17 natural populations from across three of the proposed regions of origin and of an Italian and a Hungarian population. Here also, moderate nuclear genetic diversity was observed within the populations and the differentiation between them was similar to that observed by Mijnsbrugge et al. (2013)

but higher than that found in seedstocks (Eimert et al. 2012). While Eimert et al. (2012) and Leinemann et al. (2014) analyzed genetic diversity in autochthonous seedstocks and natural populations of blackthorn in Germany, respectively, no direct comparison is available between the two sources. Therefore, in this paper, we analyzed the nuclear genetic diversity in natural stands of blackthorn that served as sources for seedstocks studied earlier (Eimert et al. 2012) and compared the two sets.

MATERIAL and METHODS

Plant material

Fully expanded leaves were collected from 40 individuals of each of the source populations which provided the seedstocks analyzed earlier (Eimert et al. 2012). That

may or may not include the actual shrubs from which those seeds were collected, as those were not labeled at that time. Populations sampled (Figure 1) were located in (at that time) two regions of origin (BMVL 2003). Populations “Hö” (near the town of Höxter in the State of North Rhine- Westphalia) and “Fu” (near the town of Fulda in the State of Hesse) belong to the designated region of origin 4, while population “Rh” (in the Rheingau Region of the State of Hesse) was located in the then region of origin 6. For samples from population sampled in wild stands “WT” is added to the label. For the corresponding seedstocks, samples of the corresponding regions of origin from earlier studies (Eimert et al. 2012) were re-analyzed. Those population are additionally labelled “Nu” (nursery material) and “ud.” in case undifferentiated material was used.




Figure 1 Schematic geographical map of Germany showing the sources of autochthonous plant material used in the recent analysis. Populations: Hö (Höxter), Fu (Fulda), Rh (Rheingau); Numbers refer to the designated regions of origin (BMELV 2003); bar 100 km

HAT-RAPD

Total DNA was extracted from frozen leaves according to Eimert et al. (2012). A highly reproducible HAT-RAPD protocol was used to analyze the DNAs of the freshly sampled natural population and to re-analyze the DNAs of the seedstocks used before with UBC primers 034, 060, 065, 096, 302, 319, 391, 727, 729, 746 and 766 under the same conditions (Eimert et al. 2012). PCR reactions were repeated at least twice to ensure reproducibility of the bands obtained.

Data analysis

BioNumerics software package (version 6.6.7; Applied Maths BVBA, Sint-Martens-Latem, Belgium) was used for band scoring and matching, individual cluster analyses (unweighted pair-group method with arithmetical averages – UPGMA and neighbor-joining method - NJ) and principal coordinate analysis (PCoA) and the resulting character table was exported and used for population analyses (raw data available upon request). Neutrality of the used markers was tested with Fst outlier methods implemented in LOSITAN (Antao et al. 2008) and Tajima's D pairwise

distance method implemented in Arlequin (Excoffier et al. 2005) using 1000 simulations. FAMD (Schlüter and Harris 2006) was used to compile descriptive statistics and to compute genetic distances for pairwise Fst values used in AMOVA and PCoA. Mantel test was conducted using the Isolation By Distance Web Service (Jensen et al. 2005). Genetic structure within and among the sampled populations was also analyzed using a Bayesian approach implemented in the software Structure (Pritchard et al. 2000). Geographic distances between populations were estimated using the Google Earth™ (Google 2009) “ruler” tool.

RESULTS

Population statistics

All 360 Individuals of the different sources were analysed by HAT-RAPD using 11 primers resulting in 390 scorable band classes. While a high number of those bands were polymorphic, the gene diversity within the populations remained rather low (Table 1). Fixed bands and private bands could be observed in every population. However, no fixed private bands could be identified.

Table 1: Population statistics

population (region of origin, source)	sample size	% polymorphic bands	fixed bands ₁	private bands ₂	fixed private bands ₃	h ₄
Fu (# 4, Nu)	40	65.90	10	1	0	0.1107
Hö (# 4, Nu)	40	58.21	9	3	0	0.1119
Rh (# 6, Nu)	40	59.23	8	2	0	0.1049
Ba (# 7, Nu)	40	72.82	4	2	0	0.1061
BB (ud. ₅ , Nu)	40	68.72	7	3	0	0.1161
Hu (ud., Nu)	40	74.62	1	4	0	0.1216
Fu (# 4, WT)	39	52.82	9	2	0	0.0993
Hö (# 4, WT)	42	67.95	6	12	0	0.1163
Rh (# 6, WT)	39	51.79	7	2	0	0.0968

1=bands which are monomorphic in a given population; 2=bands which are found exclusively in one given population; 3=bands which are found exclusively in one given population and are monomorphic in that population; 4 h=Nei's (1978) unbiased gene diversity (= measure for heterozygosity); 5 ud.=undifferentiated.

Also, it can be seen, that heterozygosity in all three wild populations is even lower than in the corresponding seedstock populations.

Using a stepwise mutation model with a 99.5% confidence intervall most of the loci used as markers in this study seem to behave selectively neutral in the wild autochthonous populations. Nevertheless,

several F_{st} outliers could be detected identifying possible candidates for balancing (7 loci) and for positive (12 loci) selection (Figure 2).

Tajima's test of selective neutrality revealed no significant deviations from a neutral model in any of the tested populations (Table 2).




Figure 2 F_{st} outliers detected by LOSITAN in autochthonous wild populations (red area – candidates for positive selection, yellow area – candidates for balancing selection).

Table 2: Tajima’s test of selective neutrality

	Fu (Nu)	Hö (Nu)	Rh (Nu)	Ba (Nu)	BB (Nu)	Hu (Nu)	Fu (WT)	Hö (WT)	Rh (WT)
Tajima’s D	-0.129	-0.039	-0.105	-0.396	-0.159	-0.188	0.075	-0.065	-0.081
P value	0.531	0.555	0.545	0.400	0.515	0.479	0.583	0.527	0.525

For Tajima’s D, values >2 and <-2 are usually considered significant deviations from 0.

Genetic Structure - Distance based methods

Clustering of all single individuals was conducted using UPGMA (Sokal and Micheneror 1958) and NJ (Saitou and Nei 1987) methods with either Jaccard (1901) or Dice (1945) coefficients. No significant differences in clustering were found with these different methods (data not shown).

While there are no larger clusters consisting uniformly of individuals of only one given population, one main clade consisted most of the individuals from sampled from the three wild stands, a second clade included most of the autochthonous seedstocks (except for Ba(Nu)) and a third one which combined most of the commercial seedstocks and the Bavarian autochthonous seedstock (Figure 3).




Figure 3 Radial cladogram showing the clustering of all screened individuals (neighbor joining, Jaccard).

The standard similarity was computed using Jaccard's similarity coefficient. Standard Jaccard distance transformation ($d=1-s$) was applied to calculate pairwise F_{st} values to assess differentiation among populations (Table 3). F_{st} values were calculated for both neutral and non-neutral markers and no significant difference was observed (data not shown).

While the differentiation among the seedstock populations was usually weak there is mostly moderate to even strong differentiation between the wild

populations and the seedstock populations and also among the wild populations themselves. Accordingly, the derived dendrogram apparently shows three larger clusters – a wider one consisting of all wild populations with the Rh(WT) population slightly removed and two tighter ones, one consisting of the autochthonous seedstocks (with the exception of Ba(Nu)) and another one consisting of the two undifferentiated commercial seedstocks and the Bavarian autochthonous seedstock (Figure 4).

Table 3: Pairwise Fst values and geographic distances

	Fu (Nu)	Hö (Nu)	Rh (Nu)	Ba (Nu)	BB (Nu)	Hu (Nu)	Fu (WT)	Hö (WT)	Rh (WT)
Fu (Nu)	---	133	137	155	n.d.	825*			
Hö (Nu)	0.1039	---	214	285	n.d.	927*			
Rh (Nu)	0.0861	0.1158	---	226	n.d.	900*			
Ba (Nu)	0.1194	0.1734	0.1146	---	n.d.	685*			
BB (Nu)	0.1003	0.1420	0.1132	0.0683	---	n.d.			
Hu (Nu)	0.1181	0.1594	0.1250	0.0464	0.0530	---			
Fu (WT)	0.2090	0.2719	0.2061	0.1547	0.1721	0.1642	---		
Hö (WT)	0.1779	0.2232	0.1910	0.1469	0.1587	0.1411	0.1384	---	
Rh (WT)	0.2288	0.2832	0.2406	0.1638	0.1879	0.1732	0.1841	0.1956	---

Above diagonal: pairwise geographical distances (km) (* estimate, n.d.–not determined); Below diagonal: Pairwise FST Values; values <0.05=no, 0.05-0.14=weak, 0.15-0.24=moderate, >0.25=strong genetic differentiation.




Figure 4 Dendrogram of the studied populations based on UPGMA analysis of pairwise distances using the Jaccard similarity index: Areas bordered by slashed lines indicate the three main clusters.

Multivariate analysis (PCoA) reveals that the wild populations are most differentiated with the Rh (WT) being the most distant (Figure 5). The wild populations are loosely grouped while the

seedstock populations group tighter and the autochthonous seedstocks group together (except Ba(Nu)) and even farther than the commercial ones.




Figure 5 Multivariate (PCoA) analysis of seedstock and wild populations

An AMOVA revealed that, in all the seedstock populations the higher genetic diversity was mostly due to variation within the populations (92.218%) and less among them (7.782%). In the wild populations, the generally lower genetic diversity is shifted to 77.473% variation within and 22.527% among populations, respectively.

However, using Fisher's exact test for population differentiation (Raymond and Rousset 1995) no differentiation between any of the analyzed populations could be detected at the 95% significance interval (after 30 000 Markov transformations).

Testing for isolation by distance (Wright 1943) a Mantel Test showed no significant correlation between genetic and geographic distances in either seedstocks ($r=0.1770$, $p=0.2200$, 999 permutations) or the wild populations ($r=0.0895$, $p=0.2330$, 999 permutations).

Genetic Structure – Bayesian Estimates

Possible genetic structures were also analyzed using the Bayesian approach based on the most likely k value (Evanno et al. 2005). Using the most sensitive settings a weak genetic structure ($\Delta K=48.33$) was shown with the most likely $k=6$ (Figure 6).




Figure 6 Bayesian clustering analysis of blackthorn populations for $K = 6$

For the highest probable $K(=6)$ all autochthonous seedstock populations (except Ba(Nu)) are located in one cluster. Of the wild populations, two (Fu(WT) and Hö(WT)) are grouped together while the third one (Rh(WT)) is farther removed and considered an extra cluster. Each of the commercial sources (BB(Nu) and

Hu(Nu)) and the remaining autochthonous seedstock (Ba(Nu)) are much more divers and each constituted a separate cluster. When we remove those last three populations from analysis the Bayesian approach detects a further genetic structuring (Figure 7), though with a much lower ΔK value (8.97).




Figure 7 Bayesian clustering analysis of autochthonous sources of blackthorn with $K=4$

With this weak structuring, all autochthonous seedstocks cluster together, while each autochthonous wild population is differentiated from them and from each other.

DISCUSSION

The above results on genetic differentiation between the tested autochthonous populations being about 22.5% of the total genetic variation are within the range reported by other authors in autochthonous blackthorn populations in Germany and bordering Flanders (Mijnsbrugge et al. 2013, Leinemann et al. 2014). In this, these results differ from the 7.8% observed earlier between the seedstocks populations harvested from

exactly the same autochthonous wild populations (Eimert et al. 2012). Also, heterozygosity observed in the wild populations was lower than that of the corresponding seedstock populations. Thus, the situation observed in the wild autochthonous populations (lower genetic diversity and higher genetic differentiation) seems to follow the common assumption about genetic adaption of (isolated) populations to the local conditions. However, we submit that the picture is not that clearcut. If the differentiation we observed were due to adaptation, one would expect a deviation from Hardy-Weinberg equilibrium. Yet, no such obvious deviation could be detected – Tajima's D shows now no significant deviation from a selective neutrality model,

and only very few loci seemed to be under selective pressure as judged from the number of *Fst* outliers. Also, the fact that no fixed private bands could be detected in 390 markers seems to contradict an IBA (isolation by adaptation) scenario. If, on the other hand, the differentiation was due to gene flow and drift assumed for the neutral theory of molecular evolution (Kimura 1968) one would expect an isolation by distance (IBD) effect for populations geographically and or ecologically separated from each other. A Mantel test on that account did not confirm such effect. Furthermore, the amount of gene flow calculated from these weak *Fst* values is accordingly high with $Nm=5.558$ ($Nm = 0.5(1 - Fst)/Fst$) meaning that to explain this low differentiation between our autochthonous populations a migration of more than 5 individuals per generation into each population would be required. As the geographical distance between those populations is more than 130 – 210 km this seems an unlikely event. On the other hand, the measurement of gene flow cannot distinguish between recent or historical events (Lowe et al. 2008). Humans have apparently used blackthorn fruits for consumption since the neolithic (Karg and Markle 2002, Martin et al. 2008) and it was cultivated in Northern Europe since the Roman Empire (Karnitsch 1953). Thus, anthropogenic distribution of blackthorn seems very plausible.

It also has been noted, that in many cases the impact of processes on genetic structuring can be obscured (Orsini et al. 2013). Thus, it can be difficult to distinguish the results of IBD and IBC (isolation by colonization). These authors propose to study the patterns of variation of both neutral and non-neutral markers in order to distinguish between the different driving forces of genetic differentiation. Using those parameters, we observe a pattern more akin to that described by

Orsini et al. (2013) as typical for IBC rather than that for IBD or IBA (isolation by adaptation) in that the *Fst* values for neither the neutral nor the non-neutral loci vary significantly over geographical distance.

Summarizing, we observe a weak pattern of differentiation between wild autochthonous populations of blackthorn in Germany. No obvious deviation from neutral evolution was detected and, thus, isolation by distance or by adaptation seems unlikely. The observed patterns support an (weak) isolation by colonization probably further driven by landscape fragmentation.

On the other hand, the situation in seedstock populations collected from those wild stands is different. That the genetic diversity within those populations is higher is not surprising by itself, considering the almost ubiquitous occurrence of blackthorn in Germany (although mostly in fragmented smaller stands), its outcrossing biology and tetraploidy. At the same time, the genetic differentiation between the seedstock populations is smaller than that between their corresponding source populations. That leads to the situation that, genetically, the wild autochthonous populations are more similar to each other than to their own more heterozygous offspring (Fig.3).

This situation presents some problems for conservation measures. If the main function of the Conservation Law is to maintain the genetic *status quo* of an existing population/sub-population the usage of residential seedstocks will not necessarily accomplish that goal in full. If the weak differentiation we observed in wild autochthonous blackthorn populations is indeed mainly due to colonization even the use of autochthonous seedstock will not maintain the same narrow genetic composition. That will

obviously depend on the concerned species, its current genetic composition and its reproductive biology. Thus, from a biological point of view an accordingly adapted approach for each species would be appropriate.

For blackthorn it seems that, at least for the regions of origin 4 and 6, the observed genetic situation does not correspond with the proposed geographic differentiation, as has been shown for other regions (Leinemann et al. 2014). A more thorough analysis of populations from all regions of Germany would be required to propose corresponding and biologically meaningful regions of origin. A recently implemented further reduction of the proposed number of regions of origin for autochthonous plants from 9 to 6 (BMU 2012) might be nearer to the “genetic truth” for blackthorn. Nonetheless, the situation will have to be evaluated for each species separately.

ACKNOWLEDGEMENTS

This work has in part been developed through funding from Hochschule RheinMain University of Applied Sciences Wiesbaden Rüsselsheim Geisenheim (grant FHW-Nr. 40130130). We are grateful to Mrs. Hüwe for excellent technical support.

REFERENCES

- Antao T, Lopes A, Lopes RJ, Beja-Pereira A, Luikart, G (2008) LOSITAN: a workbench to detect molecular adaptation based on a Fst-outlier method. *BMC Bioinformatics* 9: 323
- BMU (Bundesministerium für Umwelt, Naturschutz und Reaktorsicherheit) (2012) Leitfaden zur Verwendung gebietseigener Gehölze. Selbstverlag, Berlin
- BMVEL (Bundesministerium für Verbraucherschutz, Ernährung und Landwirtschaft) (2003) Verwendung einheimischer Gehölze regionaler Herkunft für die freie Landschaft. Ein Beitrag zur Erhaltung und Förderung der biologischen Vielfalt. Selbstverlag, Bonn
- BnatSchG (Gesetz über Naturschutz und Landschaftspflege) (2010) Bundesnaturschutzgesetz vom 29.07.2009, in Kraft getreten am 01.03.2010. BGBl I:2542
- Convention on biological diversity (1992) Secretariat of the convention on biological diversity. <http://www.cbd.int/programmes/outreach/awareness/publications.shtml>. Accessed 18th August 2011
- Crawford KM, Whitney KD (2010) Population genetic diversity influences colonization success. *Mol Ecol* 19(6):1253–1263
- Dice LR (1945) Measures of the amount of ecologic association between species. *Ecology* 26: 297-302
- Eimert K, Rückert F-E, Schröder M.B. (2011) Genetic diversity within and between seedstock populations of several German autochthonous provenances and conventionally propagated nursery material of blackthorn (*Prunus spinosa* L.). *Plant Syst Evol* 298(3):609–618
- Esquinas-Alcázar J (2005) Science and society: protecting crop genetic diversity for food security: political, ethical and technical challenges. *Nat Rev Genet* 6(12):946–953
- Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol* 14(8):2611–2620
- Excoffier L, Laval G, Schneider S (2005) Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1:47–50
- Fronia R (2009) Prüfung der Identität und Variabilität gebietsheimischer und gebietsfremder Herkünfte von *Prunus spinosa* L. und *Cornus sanguinea* L. zur Verwendung in der freien Landschaft. Der Andere Verlag, Tönning, Lübeck, Marburg
- Google (2014) Google Earth. Available: <http://www.google.com/earth>
- Gutián J, Gutián P, Sánchez JM (1993) Reproductive biology of two *Prunus* species (Rosaceae) in the Northwest Iberian Peninsula. *Plant Syst Evol* 185(3-4):153–165
- Jensen JL, Bohonak AJ, Kelley ST (2005) Isolation by distance, web service. *BMC Genet* 6(1):13
- Leinemann L, Bendixen K, Kownatzki D, Hattemer HH, Liepe K, Stenger G (2002) Genetische Untersuchungen an Landschaftsgehölzen im Hinblick auf die Erzeugung und Zertifizierung von Vermehrungsgut. *Allgemeine Forst- und Jagdzeitung* 173:146-152
- Leinemann L, Kleinschmit J, Fussi B, Hosius B, Kuchma O, Arenhövel W, Finkeldey R (2014) Genetic composition and differentiation of sloe (*Prunus spinosa* L.) populations in Germany with respect to the tracing of reproductive plant

- material. *Plant Syst Evol.* doi:10.1007/s00606-014-1027-7
- Vander Mijnsbrugge K (2013) Genetic and morphological variability among autochthonous *Prunus spinosa* populations in Flanders (northern part of Belgium): implications for seed sourcing. *Plant Ecol Evol* 146(2):193–202
- Mohanty A, Martín JP, Aguinalgalde I (2000) Chloroplast DNA diversity within and among populations of the allotetraploid *Prunus spinosa* L. *Theor Appl Genet* 100(8):1304–1310
- Mohanty A, Martín JP, Aguinalgalde I (2002) Population genetic analysis of European *Prunus spinosa* (Rosaceae) using chloroplast DNA markers. *Am J Bot* 89(8):1223–1228
- Reynders-Aloisi S, Grellet F (1994) Characterization of the ribosomal DNA units in two related *Prunus* species (*P. Cerasifera* and *P. Spinosa*). *Plant Cell Rep* 13(11):641–646
- Hoisington D, Khairallah M, Reeves T, Ribaut J-M, Skovmand B, Taba S, Warburton M (1999) Plant genetic resources: What can they contribute toward increased crop productivity? *P Natl Acad Sci USA* 96(11):5937–5943
- Jaccard P (1901) Distribution de la flore alpine dans le bassin des Dranses et dans quelques régions voisines. *Bull Soc Vaud Sci Nat* 37:241–272
- Karg S, Markle T (2002) Continuity and changes in plant resources during the Neolithic period in western Switzerland. *Veg Hist Archaeobot* 11:169–176
- Karnitsch P (1953) Fundberichte im Jahrbuch der Stadt Linz 1951. *Pro Austria Romana* 3:26
- Kimura, M (1968) Evolutionary rate at the molecular level. *Nature* 217(5129):624–626
- Kremer A, Kleinschmit J, Cottrell J, Cundall EP, Deans JD, Ducouso A, Stephan BR (2002) Is there a correlation between chloroplastic and nuclear divergence, or what are the roles of history and selection on genetic diversity in European oaks? *Forest Ecol Manag* 156(1-3):75–87
- Lowe A, Harris S, Ashton P (2008) *Ecological genetics: design, analysis, and application.* Blackwell, Malden
- Magri D, Vendramin GG, Comps B, Dupanloup I, Geburek T, Gömöry D, de Beaulieu J-L (2006) A new scenario for the quaternary history of European beech populations: palaeobotanical evidence and genetic consequences. *The New Phytologist*, 171(1), 199–221
- Martin L, Jacomet S, Thiebault S (2008) Plant economy during the Neolithic in a mountain context: the case of “Le Chenet des Pierres” in the French Alps (Bozel-Savoie, France). *Veg Hist Archaeobot* 17:113–122
- Nei M (1978) Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* 89(3):583–590.
- Orsini L, Vanoverbeke J, Swillen I, Mergeay J, De Meester L (2013) Drivers of population genetic differentiation in the wild: isolation by dispersal limitation, isolation by adaptation and isolation by colonization. *Mol Ecol* 22(24):5983–5999
- Petit RJ, Aguinalgalde I, de Beaulieu J-L, Bittkau C, Brewer S, Cheddadi R, Vendramin GG (2003) Glacial refugia: hotspots but not melting pots of genetic diversity. *Science* 300(5625):1563–1565
- Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* 155 (2):945–959.
- Raymond M, Rousset F (1995) An exact test for population differentiation. *Evolution* 49(6):1280
- Rees M, Condit R, Crawley M, Pacala S, Tilman D (2001) Long-term studies of vegetation dynamics. *Science* 293(5530):650–655
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4(4):406–425.
- Schlüter PM, Harris SA (2006) Analysis of multilocus fingerprinting data sets containing missing data. *Mol Ecol Notes* 6(2):569–572
- Schmidt PA, Krause A (1997) Zur Abgrenzung von Herkunftsgebieten bei Baumschulgehölzen für die freie Landschaft. *Natur und Landschaft* 72:92–95
- Schütt P, Schuck HJ, Stimm B (eds) (1992) *Lexikon der Forstbotanik.* Verlag Hüthig, Jehle Rehm, Heidelberg
- Sokal R and Michener C (1958) A statistical method for evaluating systematic relationships". *University of Kansas Science Bulletin* 38: 1409–1438
- Wright S (1943) Isolation by Distance. *Genetics* 28(2):114–138